

**Local Magnetization in Critical Ising Model
with Boundary Magnetic Field**

R.Chatterjee [†] and A.Zamolodchikov ^{*}
 Department of Physics and Astronomy
 Rutgers University
 P.O.Box 849, Piscataway, NJ 08855-0849

Abstract

We describe a simple way to derive spin correlation functions in 2D Ising model at critical temperature but with nonzero magnetic field at the boundary. Local magnetization (i.e. one-point function) is computed explicitly for half-plane and disk geometries.

As is known, the continuous limit of 2D Ising model at critical temperature $T = T_c$ is described by $c = 1/2$ conformal field theory which admits a Lagrangian formulation as a theory of free massless Majorana Fermi field $(\psi, \bar{\psi})$ with the action

$$\mathcal{A} = \frac{1}{2\pi} \int d^2z [\psi \partial_{\bar{z}} \psi + \bar{\psi} \partial_z \bar{\psi}]. \quad (1)$$

Here $(z, \bar{z}) = (x + iy, x - iy)$ are complex coordinates of euclidean plane and $d^2z = dx dy$. The factor $-1/2\pi$ in this action corresponds to standard normalization of the Fermi field, i.e.

$$\psi(z)\psi(z') = \frac{1}{z - z'} + \dots; \quad \bar{\psi}(\bar{z})\bar{\psi}(\bar{z}') = \frac{1}{\bar{z} - \bar{z}'} + \dots. \quad (2)$$

The order parameter field $\sigma(w, \bar{w})$ is related to the above Fermi field non-locally (it creates square-root branch cut for $\psi, \bar{\psi}$) so that the the components $\psi(z), \bar{\psi}(\bar{z})$ satisfy operator product expansions

$$\psi(z)\sigma(w, \bar{w}) = (z - w)^{-\frac{1}{2}} \sum_{n=0}^{\infty} (z - w)^n a_{-n} \sigma(w, \bar{w});$$

[†] email: robin@physics.rutgers.edu

^{*} Supported in part by Department of Energy under grant No. DE-FG0590ER40559

$$\bar{\psi}(\bar{z})\sigma(w, \bar{w}) = (\bar{z} - \bar{w})^{-\frac{1}{2}} \sum_{n=0}^{\infty} (\bar{z} - \bar{w})^n \bar{a}_{-n} \sigma(w, \bar{w}), \quad (3)$$

where a_{-n} and $\bar{a}_{-n}$ are mode operators associated with ψ and $\bar{\psi}$. The field $a_0\sigma$ is known as “disorder parameter” field μ ; in fact we have

$$a_0\sigma(w, \bar{w}) = \frac{\omega}{\sqrt{2}}\mu(w, \bar{w}); \quad \bar{a}_0\sigma(w, \bar{w}) = \frac{\bar{\omega}}{\sqrt{2}}\mu(w, \bar{w}), \quad (4)$$

where

$$\omega = e^{\frac{i\pi}{4}}; \quad \bar{\omega} = e^{-\frac{i\pi}{4}}, \quad (5)$$

and we have assumed the standard normalization of the fields σ and μ , i.e.

$$\sigma(w, \bar{w})\sigma(0, 0) = (w\bar{w})^{-\frac{1}{8}} \times I + \dots; \quad \mu(w, \bar{w})\mu(0, 0) = (w\bar{w})^{-\frac{1}{8}} \times I + \dots, \quad (6)$$

where I is the identity operator. In fact, the first two subleading terms in (3) are expressed in terms of derivatives of μ , namely

$$a_{-1}\sigma(w, \bar{w}) = \frac{\omega}{\sqrt{2}}(4\partial_w\mu(w, \bar{w})); \quad a_{-2}\sigma(w, \bar{w}) = \frac{\omega}{\sqrt{2}}(\frac{8}{3}\partial_w^2\mu(w, \bar{w})). \quad (7)$$

$$\bar{a}_{-1}\sigma(w, \bar{w}) = \frac{\bar{\omega}}{\sqrt{2}}(4\partial_{\bar{w}}\mu(w, \bar{w})); \quad \bar{a}_{-2}\sigma(w, \bar{w}) = \frac{\bar{\omega}}{\sqrt{2}}(\frac{8}{3}\partial_{\bar{w}}^2\mu(w, \bar{w})). \quad (8)$$

These equations are easily obtained with the use of explicit expressions

$$T(z) = -\frac{1}{2}(\psi\partial_z\psi)(z); \quad \bar{T}(\bar{z}) = -\frac{1}{2}(\bar{\psi}\partial_{\bar{z}}\bar{\psi})(\bar{z}) \quad (9)$$

for the components $T = T_{zz}$ and $\bar{T} = T_{\bar{z}\bar{z}}$ of stress-energy tensor in terms of Fermi fields $\psi, \bar{\psi}$. Similar equations hold for the products $\psi\mu, \bar{\psi}\mu$;

$$\begin{aligned} \psi(z)\mu(w, \bar{w}) &= \frac{\bar{\omega}}{\sqrt{2}}(z - w)^{-\frac{1}{2}}[\sigma(w, \bar{w}) + 4(z - w)\partial_w\sigma(w, \bar{w}) + \frac{8}{3}(z - w)^2\partial_w^2\sigma(w, \bar{w}) + \dots]; \\ \bar{\psi}(\bar{z})\mu(w, \bar{w}) &= \frac{\omega}{\sqrt{2}}(\bar{z} - \bar{w})^{-\frac{1}{2}}[\sigma(w, \bar{w}) + 4(\bar{z} - \bar{w})\partial_{\bar{w}}\sigma(w, \bar{w}) + \frac{8}{3}(\bar{z} - \bar{w})^2\partial_{\bar{w}}^2\sigma(w, \bar{w}) + \dots]. \end{aligned} \quad (10)$$

This form of operator product expansions makes it possible to derive linear differential equations which determine multipoint correlation functions of the fields σ and μ (see e.g. [1]).

Conformal field theory of critical Ising model in the presence of boundary (along with more general conformal field theories) is studied in Refs[2-4]. It is shown in [3] that there are two essentially different possibilities to choose “critical” boundary conditions, the ones which preserve conformal invariance. If the boundary $\mathcal{B}$ is given by a parametric curve

$$\mathcal{B}: \quad z = Z(t); \quad \bar{z} = \bar{Z}(t), \quad (11)$$

where t is some real parameter along the boundary, these two possibilities are

$$[e^{\frac{1}{2}}\psi - \bar{e}^{\frac{1}{2}}\bar{\psi}]_{\mathcal{B}} = 0 \quad (12)$$

and

$$[e^{\frac{1}{2}}\psi + \bar{e}^{\frac{1}{2}}\bar{\psi}]_{\mathcal{B}} = 0, \quad (13)$$

where

$$e(t) = \frac{d}{dt}Z(t), \quad \bar{e}(t) = \frac{d}{dt}\bar{Z}(t) \quad (14)$$

are the components of the vector $(e, \bar{e})$ tangent to the boundary; in what follows we assume that the parametrization (11) of the curve $\mathcal{B}$ is chosen in such a way that this vector has unit length, i.e.

$$e(t)\bar{e}(t) = 1; \quad (15)$$

we also assume the choice of orientation of this curve in which the unit vector $(ie, -i\bar{e})$ normal to the boundary points inside the domain $\mathcal{D}$ surrounded by $\mathcal{B}$; $\mathcal{B} = \partial\mathcal{D}$. From a microscopic point of view, (12) and (13) correspond to “free” and “fixed” boundary conditions, respectively. Namely, (12) appears when the microscopic boundary spins are not restricted, while (13) corresponds to the situation when the boundary spins are fixed to be all in the same position, $+1$ or -1 (so that (13) represents, in fact, two boundary conditions, “fixed to $+1$ ” and “fixed to -1 ”). In both cases, due to the conformal invariance, one can reduce the case of a generic (one-component) curve $\mathcal{B}$ to the “standard” geometry where $\mathcal{D}$ is the upper half-plane $y > 0$. Then (12) and (13) simplify as

$$[\psi - \bar{\psi}]_{z=\bar{z}} = 0 \quad \text{for “free” case}, \quad (12a)$$

$$[\psi + \bar{\psi}]_{z=\bar{z}} = 0 \quad \text{for “fixed” case} \quad (13a)$$

These equations show that the field $\bar{\psi}(\bar{z})$ coincides (up to a sign which depends on whether we choose “free” or “fixed” boundary condition) with the analytic continuation of $\psi(z)$ to the lower half-plane. Therefore it is still possible to use (7,8,10) to derive complete set of linear differential equations which determine correlation functions of the fields σ and μ in the presence of boundary [2].

The above “free” and “fixed” boundary conditions are “critical”, i.e. they correspond to “fixed points” of the renormalization group flow of “boundary interactions”, and associated field theories with these boundary conditions are conformally invariant. In this paper we consider more general boundary conditions which break conformal symmetry. Specifically, we analyze the case when boundary spins interact with constant external magnetic field H_B . Clearly, this boundary condition “interpolates” between the “free”

case at $H_B = 0$ and “fixed to ± 1 ” cases at $H_B \rightarrow \pm\infty$, and so it corresponds to “boundary flow” from “free” boundary condition down to “fixed” one[5]. This “flow” can be interpreted as “free” boundary condition “perturbed” by “boundary spin operator” $\sigma_B(t)$. The latter was identified in [3,4] (see also [6]); it can be expressed in terms of Fermi fields as

$$\sigma_B(t) = i a(t) [e^{\frac{1}{2}} \psi + \bar{e}^{\frac{1}{2}} \bar{\psi}](t), \quad (16)$$

where $a(t)$ is auxiliary “boundary” Fermionic degree of freedom with the two-point function $\langle a(t)a(t') \rangle_{free} = \frac{1}{2} \text{sign}(t-t')$ [6]. With this, the full action describing the continuous limit of $T = T_c$ Ising model with boundary magnetic field takes the form

$$\mathcal{A}_h = \frac{1}{2\pi} \int_{\mathcal{D}} d^2z [\psi \partial_{\bar{z}} \psi + \bar{\psi} \partial_z \bar{\psi}] + \int_{\mathcal{B}} [-\frac{i}{4\pi} \psi \bar{\psi} + \frac{1}{2} a \dot{a}] dt + i h \int_{\mathcal{B}} a(t) (e^{\frac{1}{2}} \psi + \bar{e}^{\frac{1}{2}} \bar{\psi})(t) dt, \quad (17)$$

where $\dot{a} \equiv \frac{d}{dt} a$. The first two terms here give the action for Ising model field theory with “free” boundary condition (10) and the last term is the “perturbation” describing the interaction with the boundary field; it contains dimensional constant $h \sim [length]^{-1}$ (appropriately rescaled external field H_B) and breaks conformal symmetry. Note that the full action is quadratic, so it is still a free field theory. The fact that the boundary magnetic field does not destroy free-field structure of Ising model is very well known (see e.g.[7]). Nevertheless, we consider it interesting to observe that although conformal symmetry is broken the correlation functions still satisfy linear differential equations.

The boundary condition for the fields $\psi, \bar{\psi}$ in the presence of the boundary field h is derived directly from (17),

$$(\frac{d}{dt} + i\lambda)\psi(t) = (\frac{d}{dt} - i\lambda)\bar{\psi}(t), \quad (18)$$

where

$$\psi(t) = e^{\frac{1}{2}}(t) \psi(Z(t)); \quad \bar{\psi}(t) = \bar{e}^{\frac{1}{2}}(t) \bar{\psi}(Z(t)), \quad (19)$$

and

$$\lambda = 4\pi h^2. \quad (20)$$

We want to show that in the presence of boundary magnetic field h the correlation functions of σ and μ still satisfy linear differential equations. We will demonstrate this explicitly for the one-point function $\langle \sigma(z, \bar{z}) \rangle_h$ (“local magnetization”) in the simplest case of half-plane geometry. So, we assume that the boundary $\mathcal{B}$ is the real axis $z = \bar{z}$ and $\mathcal{D}$ is the upper half-plane $y > 0$. In this case the boundary condition (19) reduces to

$$[(\partial_z + i\lambda)\psi(z) - (\partial_{\bar{z}} - i\lambda)\bar{\psi}(\bar{z})]_{z=\bar{z}} = 0. \quad (21)$$

This form of the boundary condition makes it explicit that the fields

$$\chi(z) = (\partial_z + i\lambda)\psi(z); \quad \bar{\chi}(\bar{z}) = (\partial_{\bar{z}} - i\lambda)\bar{\psi}(\bar{z}), \quad (22)$$

enjoy the desired property that $\bar{\chi}(\bar{z})$ coincides with the analytic continuation of $\chi(z)$ to the lower half-plane. So, for instance, the correlation function $\langle \chi(z)\mu(w, \bar{w}) \rangle$ is an analytic function of z on the full z -plane with two square-root branch points at $z = w$ and $z = \bar{w}$. Taking into account (10) and the asymptotic behavior

$$\chi(z) \sim z^{-1} \quad \text{as } z \rightarrow \infty \quad (23)$$

one can write

$$\langle \chi(z)\mu(w, \bar{w}) \rangle_h (z - w)^{\frac{1}{2}} (z - \bar{w})^{\frac{1}{2}} = \frac{A(w, \bar{w})}{z - w} + \frac{\bar{A}(w, \bar{w})}{z - \bar{w}} + B(w, \bar{w}). \quad (24)$$

Now, one can use the operator product expansions

$$\begin{aligned} \chi(z)\mu(w, \bar{w}) &= \frac{\bar{w}}{\sqrt{2}}(z - w)^{-\frac{3}{2}} \left(-\frac{1}{2}\sigma(w, \bar{w}) + (z - w)(2\partial_w + i\lambda)\sigma(w, \bar{w}) + \right. \\ &\quad \left. + (z - w)^2(4\partial_w^2 + 4i\lambda\partial_w)\sigma(w, \bar{w}) + \dots \right); \end{aligned} \quad (25)$$

and

$$\begin{aligned} \bar{\chi}(\bar{z})\mu(w, \bar{w}) &= \frac{\omega}{\sqrt{2}}(\bar{z} - \bar{w})^{-\frac{3}{2}} \left(-\frac{1}{2}\sigma(w, \bar{w}) + (\bar{z} - \bar{w})(2\partial_{\bar{w}} - i\lambda)\sigma(w, \bar{w}) + \right. \\ &\quad \left. + (\bar{z} - \bar{w})^2(4\partial_{\bar{w}}^2 - 4i\lambda\partial_{\bar{w}})\sigma(w, \bar{w}) + \dots \right), \end{aligned} \quad (26)$$

which follow from (10), to express the coefficients $A, \bar{A}$ and B in (24) in terms of the function $\langle \sigma(w, \bar{w}) \rangle$ and its derivatives. For instance, expanding (24) in powers of $z - w$ with the use of (25) we get

$$A(w, \bar{w}) = \frac{\bar{w}}{\sqrt{2}}(w - \bar{w})^{\frac{1}{2}} \left(-\frac{1}{2} \right) \langle \sigma(w, \bar{w}) \rangle_h; \quad (27a)$$

$$B(w, \bar{w}) + \frac{\bar{A}(w, \bar{w})}{w - \bar{w}} = \frac{\bar{w}}{\sqrt{2}}(w - \bar{w})^{\frac{1}{2}} \left(2\partial_w + i\lambda - \frac{1}{4} \frac{1}{w - \bar{w}} \right) \langle \sigma(w, \bar{w}) \rangle_h; \quad (27b)$$

$$\begin{aligned} -\frac{\bar{A}(w, \bar{w})}{(w - \bar{w})^2} &= \frac{\bar{w}}{\sqrt{2}}(w - \bar{w})^{\frac{1}{2}} \left(4\partial_w^2 + (4i\lambda + \frac{1}{w - \bar{w}})\partial_w + \right. \\ &\quad \left. + \frac{1}{2} \frac{i\lambda}{w - \bar{w}} + \frac{1}{16} \frac{1}{(w - \bar{w})^2} \right) \langle \sigma(w, \bar{w}) \rangle_h. \end{aligned} \quad (27c)$$

On the other hand, in view of (21), the expansion of this function in powers of $z - \bar{w}$ is controlled by the operator product expansion (26). Therefore we have also

$$\bar{A}(w, \bar{w}) = \frac{\omega}{\sqrt{2}}(\bar{w} - w)^{\frac{1}{2}} \left(-\frac{1}{2} \right) \langle \sigma(w, \bar{w}) \rangle_h; \quad (28a)$$

$$B(w, \bar{w}) + \frac{A(w, \bar{w})}{\bar{w} - w} = \frac{\omega}{\sqrt{2}}(\bar{w} - w)^{\frac{1}{2}} \left(2\partial_{\bar{w}} - i\lambda - \frac{1}{4} \frac{1}{\bar{w} - w} \right) \langle \sigma(w, \bar{w}) \rangle_h; \quad (28b)$$

$$\begin{aligned}
-\frac{A(w, \bar{w})}{(\bar{w} - w)^2} &= \frac{\omega}{\sqrt{2}}(\bar{w} - w)^{\frac{1}{2}}(4\partial_{\bar{w}}^2 + (-4i\lambda + \frac{1}{\bar{w} - w})\partial_{\bar{w}} - \\
&\quad - \frac{1}{2}\frac{i\lambda}{\bar{w} - w} + \frac{1}{16}\frac{1}{(\bar{w} - w)^2})\langle\sigma(w, \bar{w})\rangle_h.
\end{aligned} \tag{28c}$$

Compatibility of (27a,b) and (28a,b) requires that

$$(\partial_w + \partial_{\bar{w}})\langle\sigma(w, \bar{w})\rangle_h = 0. \tag{29}$$

This equation just expresses the translational symmetry of the system: the one-point function $\langle\sigma(w, \bar{w})\rangle_h$ does not depend on $x = \frac{1}{2}(w + \bar{w})$. With this, (28c) becomes identical to (27c) and reduces to the ordinary second-order differential equation

$$[4\frac{d^2}{dY^2} + (\frac{1}{Y} - 4)\frac{d}{dY} + (-\frac{1}{Y} + \frac{9}{16}\frac{1}{Y^2})]\bar{\sigma}(Y) = 0 \tag{30}$$

for the function

$$\bar{\sigma}(2\lambda y) = \langle\sigma(iy, -iy)\rangle_h. \tag{31}$$

This can be reduced to degenerate hypergeometric equation and we find that the only solution which does not grow exponentially as $Y \rightarrow +\infty$ is

$$\bar{\sigma}(Y) = C \times Y^{\frac{3}{8}}\Psi(1/2, 1, Y), \tag{32}$$

where

$$\Psi(a, c, x) = \frac{1}{\Gamma(a)} \int_0^\infty e^{-xt} t^{a-1} (1+t)^{c-a-1} dt. \tag{33}$$

The Y -independent constant C can be fixed by comparing the $Y \rightarrow 0$ asymptotic of the solution (32)

$$\bar{\sigma}(Y) = C \times \frac{1}{\sqrt{\pi}} Y^{\frac{3}{8}} [-\log(Y) + O(1)] \tag{34}$$

with the first nontrivial perturbative contribution

$$\begin{aligned}
\langle\sigma(iy, -iy)\rangle_h &= h \int_{-L}^{+L} \langle\sigma(iy, -iy)\sigma_B(x)\rangle_{free} dx + O(h^3) = \\
&= 2^{\frac{5}{4}} h (2y)^{\frac{3}{8}} (\log(2L) - \log(2y)) + O(h^3);
\end{aligned} \tag{35}$$

where we introduced an infrared cut-off L and assumed that the field σ is normalized as in (6). Finally,

$$\langle\sigma(iy, -iy)\rangle_h = \lambda^{\frac{1}{2}} 2^{\frac{1}{4}} (2y)^{\frac{3}{8}} \Psi(1/2, 1, 2\lambda y). \tag{36}$$

Note that the $Y \rightarrow \infty$ asymptotic

$$\langle \sigma(iy, -iy) \rangle_h \sim 2^{\frac{1}{4}} (2y)^{-\frac{1}{8}} \quad (37)$$

of this function coincides with $\langle \sigma(iy, -iy) \rangle_{fixed}$ (see [4]), as it should, since $h \rightarrow \infty$ limit corresponds to the “fixed to +1” boundary condition.

The above analysis can be repeated in the case when $\mathcal{D}$ is a disk of radius R , i.e. the boundary $\mathcal{B}$ is the circle

$$Z(t) = R \exp(\frac{it}{2\pi R}); \quad \bar{Z}(t) = R \exp(-\frac{it}{2\pi R}). \quad (38)$$

Obviously, in this case the one-point function $\langle \sigma(z, \bar{z}) \rangle_h$ depends only on the combination $z\bar{z}$, i.e.

$$\langle \sigma(z, \bar{z}) \rangle_h = s(z\bar{z}/R^2), \quad (39)$$

and the function $s(X)$ can be shown to satisfy

$$[4\frac{d^2}{dX^2} + (\frac{4(1+g)}{X} - \frac{1}{1-X})\frac{d}{dX} + (\frac{9}{16}\frac{1}{(1-X)^2} + \frac{1-g}{2X(1-X)})]s(X) = 0, \quad (40)$$

where $g = \lambda R$. This equation is solved in terms of hypergeometric functions; the solution which is regular everywhere inside the disk $\mathcal{D}$ is

$$\langle \sigma(z, \bar{z}) \rangle_h = g^{\frac{1}{2}} \frac{\Gamma(1/2 + g)}{\Gamma(1 + g)} (R/4)^{-\frac{1}{8}} (1 - X)^{\frac{3}{8}} F(1/2, 1/2 + g, 1 + g, X), \quad (41)$$

where

$$X = \frac{z\bar{z}}{R^2}, \quad g = 4\pi h^2 R, \quad (42)$$

and $F(a, b, c, X)$ is the hypergeometric function. Again, we used the first nontrivial order of perturbation theory in h to fix the overall normalization factor in (41). In the limit when R goes to infinity while the distance from $(z, \bar{z})$ to $\mathcal{B}$ is kept finite (i.e. $X \rightarrow 1$ with $Y = \lambda R(1 - X)$ fixed), (41) degenerates to (32).

Similar arguments can be used to derive differential equations for multipoint functions. However, these equations are more complicated and we do not describe them here.

Acknowledgement

RC would like to thank V.Brazhnikov and Y.Pugay for helpful discussions.

References

1. “Conformal Invariance and Applications to Statistical Mechanics”, C.Itzykson, H.Saleur, J.-B.Zuber, eds., World Scientific, 1988.
- 2.J.Cardy. “Conformal Invariance and Surface Critical Behavior”, in Ref.1.
- 3.J.Cardy. Nucl.Phys. B324 (1989) p.581.
- 4.J.Cardy, D.Lewellen. Phys.Lett. 259B (1991) p.274.
- 5.I.Affleck, A.Ludwig. “Universal Non-Integer “Groundstate Degeneracy” in Critical Quantum Systems”, Preprint UBCTP-91-007.
- 6.S.Ghoshal, A.Zamolodchikov. “Boundary S-matrix and Boundary State in Two-Dimensional Integrable Quantum Field Theory”. Preprint RU-93-20.
- 7.B.M.McCoy, T.T.Wu. “The Two-Dimensional Ising Model”, Harvard University Press, 1973.